

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
5 July 2001 (05.07.2001)

PCT

(10) International Publication Number
WO 01/48183 A2

- (51) International Patent Classification⁷: C12N 15/00 (74) Agent: BAYLISS, Geoffrey, Cyril; Boulton Wade Tennant, Verulam Gardens, 70 Gray's Inn Road, London WC1X 8BT (GB).
- (21) International Application Number: PCT/EP00/13149
- (22) International Filing Date: 22 December 2000 (22.12.2000) (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 9930691.2 24 December 1999 (24.12.1999) GB (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): DEV-GEN NV [BE/BE]; Technologiepark 9, B-9052 Zwijnaarde (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): PLAETINCK, Geert [BE/BE]; Pontstraat 16, B-9820 Merelbeke (BE). MORTIER, Katherine [BE/BE]; Paddenhoek 20, B-9830 St.-Martens Latem (BE). LISSENS, Ann [BE/BE]; Tiensesteenweg 137, B-3010 Kessel-Lo (BE). BOGAERT, Thierry [BE/BE]; Wolvendreef 26g, B-8500 Kortrijk (BE).
- Published:
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/48183 A2

(54) Title: IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

(57) Abstract: There are described ways of improving the efficiency of double stranded RNA inhibition as a method of inhibiting gene expression in nematode worms such as *C. elegans*. In particular, the invention relates to the finding that changes in the genetic background of *C. elegans* result in increased sensitivity to double-stranded RNA inhibition.

NUC 37139

IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNAINHIBITION

5 The present invention is concerned with ways of
improving the efficiency of double stranded RNA
inhibition as a method of inhibiting gene expression
in nematode worms such as *C. elegans*. In particular,
the invention relates to the finding that the
susceptibility of nematode worms such as *C. elegans* to
10 double stranded RNA inhibition is affected by changes
in the genetic background of the worms.

It has recently been described in Nature Vol 391,
pp.806-811, February 98, that introducing double
stranded RNA into a cell results in potent and
15 specific interference with expression of endogenous
genes in the cell, which interference is substantially
more effective than providing either RNA strand
individually as proposed in antisense technology. This
specific reduction of the activity
20 of the gene was also found to occur in the nematode
worm *Caenorhabditis elegans* (*C. elegans*) when the RNA
was introduced into the genome or body cavity of the
worm.

The present inventors have utilized the double
25 stranded RNA inhibition technique and applied it
further to devise novel and inventive methods of (i)
assigning functions to genes or DNA fragments which
have been sequenced in various projects, such as, for
example, the human genome project and which have yet
30 to be accorded a particular function, and (ii)
identifying DNA responsible for conferring a
particular phenotype. Such methods are described in
the applicant's co-pending application number WO
00/01846. Processes for introducing RNA into a living
35 cell, either *in vivo* or *ex vivo*, in order to inhibit
expression of a target gene in that cell are

CONFIRMATION COPY

NUC 37140

additionally described in WO 99/32619.

Several different experimental approaches can be used to introduce double-stranded RNA into nematode worms in order to achieve RNA interference *in vivo*.

5 One of the most straightforward approaches is simple injection of double-stranded RNA into a body cavity. A more elegant solution is to feed the nematodes on food organisms, generally bacteria, which express a double stranded RNA of the appropriate sequence,
10 corresponding to a region of the target gene.

The present inventors have now determined that the phenomenon of RNA interference in nematodes following ingestion of food organisms capable of expressing double-stranded RNA is dependent both on
15 the nature of the food organism and on the genetic background of the nematodes themselves. These findings may be exploited to provided improved methods of double-stranded RNA inhibition.

Therefore, according to a first aspect of the
20 present invention there is provided a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence
25 substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild
30 type.

Caenorhabditis elegans is the preferred nematode worm for use in the method of the invention although the method could be carried out with other nematodes and in particular with other microscopic nematodes,
35 preferably microscopic nematodes belonging to the genus *Caenorhabditis*. As used herein the term "microscopic" nematode encompasses nematodes of

approximately the same size as *C. elegans*, being of the order 1mm long in the adult stage. Microscopic nematodes of this approximate size can easily be grown in the wells of a multi-well plate of the type
5 generally used in the art to perform mid- to high-throughput screening.

It is an essential feature of this aspect of the invention that the nematode has a non wild-type genetic background which confers greater sensitivity
10 to RNA interference phenomena (abbreviated herein to RNAi) as compared to the equivalent wild type nematodes. As illustrated in the accompanying examples, introduction of double-stranded RNA (abbreviated herein to dsRNA) into a non wild-type
15 strain according to the invention results in greater inhibition of expression of the target gene. Depending on the nature of the target gene, this greater level of inhibition may be detectable at the phenotypic level as a more pronounced phenotype.

20 The nematode having non wild-type genetic background may, advantageously, be a mutant strain. Mutations which have the effect of increasing susceptibility of the nematode to RNAi may, for example, affect the stability of dsRNA or the kinetics
25 of dsRNA turnover within cells of the worm or the rate of uptake of dsRNA synthesised by a food organism. Suitable mutant strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding proteins involved in RNA
30 synthesis, RNA degradation or the regulation of these processes.

In one preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits reduced activity of one or more
35 nucleases compared to wild-type. Suitable strains include mutant strains exhibiting knock-out or loss-of-function mutations in one or more genes encoding

nucleases, such as RNases. A particularly preferred example is the *nuc-1* strain. This mutant *C. elegans* strain is known per se in the art.

5 In a second preferred embodiment, the nematode is a mutant strain, more preferably a mutant *C. elegans*, which exhibits increased gut uptake compared to wild-type. Particularly preferred examples of such strains are the so-called *C. elegans* gun mutants described herein. In a still further embodiment, the nematode
10 may be a transgenic worm comprising one or more transgenes which increase gut uptake relative to wild-type.

The term "increased gut uptake" as used herein is taken to mean increased uptake of foreign particles
15 from the gut lumen and may encompass both increased gut permeability and increased gut molecular transport compared to wild-type *C. elegans*.

C. elegans feeds by taking in liquid containing its food (e.g. bacteria). It then spits out the
20 liquid, crushes the food particles and internalises them into the gut lumen. This process is performed by the muscles of the pharynx. The process of taking up liquid and subsequently spitting it out is called pharyngeal pumping. Once the food particles have been
25 internalised via pharyngeal pumping their contents must cross the gut itself in order to reach target sites in the worm. There are multiple factors which effect the uptake of compounds from the gut lumen to the surrounding tissues. These include the action of
30 multi-drug resistance proteins, multi-drug resistance related proteins and the P450 cytochromes as well as other enzymes and mechanisms available for transport of molecules through the gut wall.

C. elegans mutants which exhibit increased uptake
35 of foreign molecules through the gut may be obtained from the *C. elegans* mutant collection at the C.

C. elegans Genetic Center, University of Minnesota, St Paul, Minnesota, or may be generated by standard methods. Such methods are described by Anderson in Methods in Cell Biology, Vol 48, "C. elegans: Modern biological analysis of an organism" Pages 31 to 58. Several selection rounds of the PCR technique can be performed to select a mutant worm with a deletion in a desired gene. Alternatively, a population of worms could be subjected to random mutagenesis and worms exhibiting the desired characteristic of increased gut uptake selected using a phenotypic screen, such as the dye uptake method described herein.

As an alternative to mutation, transgenic worms may be generated with the appropriate characteristics. Methods of preparing transgenic worms are well known in the art and are particularly described by Craig Mello and Andrew Fire, Methods in Cell Biology, Vol 48, Ed. H.F. Epstein and D.C. Shakes, Academic Press, pages 452-480.

Worms exhibiting the desired characteristics of increased gut uptake can be identified using a test devised by the inventors based on uptake of a marker precursor molecule which is cleaved by the action of enzymes present in the gut lumen to generate a marker molecule which produces a detectable signal, such as fluorescence. A suitable marker precursor molecule is the fluorescent dye precursor BCECF-AM available from Molecular Probes (Europe BV), Netherlands. This dye only becomes fluorescent when cleaved by esterases and maintained at a pH above 6. The pH of the gut lumen is usually 5 or below. Thus, any BCECF-AM taken up through the pharynx into the gut lumen is not fluorescent until cleaved and the cleaved portion has entered the cells surrounding the lumen which are at a higher pH. Thus, this dye is able to quickly identify mutant or otherwise modified worms which have increased gut transport or permeability. There is a

gradual increase in fluorescence in the tissues surrounding the gut while the gut lumen remains dark. The fluorescence can be detected at an excitation wavelength of 485 nm and an emission wavelength of 530 nm.

Specific examples of gun mutant strains isolated using this procedure which may be used in the method of the invention are strains bg77, bg84, bg85 and bg86, although it is to be understood that the invention is in no way limited to the use of these specific strains. The *C. elegans* gun mutant strain bg85 was deposited on 23 December 1999 at the BCCM/LMG culture collection, Laboratorium Voor Microbiologie, Universiteit Gent, K. L. Ledeganckstraat 35, B-9000, Gent, Belgium under accession number LMBP 5334CB. The phrase "the bg85 mutation" as used herein refers to the specific mutation(s) present in the bg85 strain which is/are responsible for conferring the gun phenotype.

It is also within the scope of the invention to use a non wild-type nematode strain, preferable a *C. elegans* strain, having multiple mutations which affect sensitivity to RNAi. A preferred type of multiple mutant is one having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type. An example of such a mutant is a *C. elegans* strain having the *nuc-1* mutation and at least one further gun mutation. As exemplified herein, double mutants having the *nuc-1* mutation and a gun mutation exhibit enhanced sensitivity to RNAi as compared to either *nuc-1* or gun single mutants.

For the avoidance of doubt, where particular characteristics or properties of nematode worms are described herein by relative terms such as "enhanced"

or "increased" or "decreased" this should be taken to mean enhanced, increased or decreased relative to wild-type nematodes. In the case of *C. elegans*, wild-type is defined as the N2 Bristol strain which is well known to workers in the *C. elegans* field and has been extremely well characterised (see Methods in Cell Biology, Volume 48, *Caenorhabditis elegans*: Modern biological analysis of an organism, ed. by Henry F. Epstein and Diane C. Shakes, 1995 Academic Press; The nematode *Caenorhabditis elegans*, ed. by William Wood and the community of *C. elegans* researchers., 1988, Cold Spring Harbor Laboratory Press; *C. elegans* II, ed. by Donald L. Riddle, Thomas Blumenthal, Barbara J. Meyer and James R. Priess, 1997, Cold Spring Harbor Laboratory Press). The N2 strain can be obtained from the *C. elegans* Genetic Center, University of Minnesota, St Paul, Minnesota, USA.

The food organism for use in the above aspect of the invention is preferably a bacterium such as, for example, a strain of *E.coli*. It will, however, be appreciated that any other type of food organism on which nematodes feed and which is capable of producing dsRNA could be used. The food organism may be genetically modified to express a double-stranded RNA of the appropriate sequence, as will be understood with reference to the examples included herein. One convenient way in which this may be achieved in a bacterial food organism is by transforming the bacterium with a vector comprising a promoter or promoters positioned to drive transcription of a DNA sequence to RNA capable of forming a double-stranded structure. Examples of such vectors will be further described below.

The actual step of feeding the food organism to the nematode may be carried out according to procedures known in the art, see WO 00/01846.

Typically the feeding of the food organisms to the nematodes is performed on standard agar plates commonly used for culturing *C. elegans* in the laboratory. However, the step of feeding the food
5 organism to the nematodes may also be carried out in liquid culture, for example in the wells of 96-well microtitre assay plates.

The inventors have further observed that variations in the food organism can result in enhanced
10 in vivo RNAi when the food organism is ingested by a nematode worm.

Accordingly, in a further aspect the invention provides a method of inhibiting expression of a target gene in a nematode worm comprising feeding to said
15 nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a
20 modification selected to provide increased expression or persistence of the doubled-stranded RNA compared to a food organism which does not carry the modification.

The modification present in the food organism can be any modification which results in increased
25 expression of the dsRNA or in increased persistence of the dsRNA. Suitable modifications might include mutations within the bacterial chromosome which affect RNA stability and/or degradation or mutations which have a direct effect on the rate of transcription. In
30 a preferred embodiment, the food organism is an RNase III minus *E. coli* strain, or any other RNase negative strain.

According to a still further aspect of the invention there is provided a method of inhibiting
35 expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide

sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

5 In addition to exhibiting increased sensitivity to RNAi following feeding with food organisms capable of expressing a dsRNA, nematodes which exhibit increase gut uptake as described herein also show increased uptake of DNA molecules capable of producing
10 double-stranded RNA structures following ingestion into a nematode.

In a preferred embodiment, the DNA is in the form of a vector comprising a promoter or promoters orientated to relative to a sequence of DNA such that
15 they are capable of driving transcription of the said DNA to make RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to the promoter or promoters.

Several different arrangements of promoters may
20 be used in such a vector. In a first arrangement a DNA fragment corresponding to a region of the target gene is flanked by two opposable polymerase-specific promoters which are preferably identical. Transcription from the opposable promoters produces
25 two complementary RNA strands which can anneal to form an RNA duplex. The plasmid pGN1 described herein is an example of a vector comprising two opposable T7 promoters flanking a multiple cloning site for insertion of a DNA fragment of the appropriate
30 sequence, corresponding to a region of a target gene. pGN8 is an example of a vector derived from pGN1 containing a fragment of the *C. elegans unc-22* gene. In an alternative arrangement, DNA fragments corresponding to a region of the target gene may be
35 placed both in the sense and the antisense orientation downstream of a single promoter. In this case, the sense/antisense fragments are co-transcribed to

generate a single RNA strand which is self-complementary and can therefore form an RNA duplex.

In both of the above arrangements, the polymerase-specific T3, T7 and SP6 promoters, all of which are well known in the art, are useful for driving transcription of the RNA. Expression from these promoters is dependent on expression of the cognate polymerase. Advantageously, the nematode itself may be adapted to express the appropriate polymerase. Expression of the polymerase may be general and constitutive, but could also be regulated under a tissue-specific promoter, an inducible promoter, a temporally regulated promoter or a promoter having a combination of such characteristics. Transgenic *C. elegans* strains harboring a transgene encoding the desired polymerase under the control of an appropriately-regulated promoter can be constructed according to methods known *per se* in the art and described, for example, by Craig Mello and Andrew Fire in *Methods in Cell Biology*, Vol 48, Ed. H. F. Epstein and D. C. Shakes, Academic Press, pp 452-480.

The advantage of adapting the nematode to express the required polymerase is that it is possible to control inhibition of expression of the target gene in a tissue-specific and/or temporally specific manner by placing expression of the polymerase under the control of an appropriately regulated promoter.

Introduction of DNA into nematodes in accordance with the method of the invention can be achieved using a variety of techniques, for example by direct injection into a body cavity or by soaking the worms in a solution containing the DNA. If the DNA is in the form of a vector as described herein, e.g. a plasmid harboring a cloned DNA fragment between two flanking T7 promoters, then dsRNA corresponding to this DNA fragment will be formed in the nematode resulting in down regulation of the corresponding gene. The

introduced DNA can form an extrachromosomal array,
which array might result in a more catalytic knock-out
or reduction of function phenotype. The DNA might also
become integrated into the genome of the nematode,
5 resulting in the same catalytic knock out or reduction
of function phenotype, but which is stably
transmittable.

In each aspect of the invention, the double-
stranded RNA structure may be formed by two separate
10 complementary RNA strands or a single self-
complementary strand, as described above. Inhibition
of target gene expression is sequence-specific in that
only nucleotide sequences corresponding to the duplex
region of the dsRNA structure are targeted for
15 inhibition.

It is preferred to use dsRNA comprising a
nucleotide sequence identical to a portion of the
target gene, although RNA sequences with minor
variations such as insertions, deletions and single
20 base substitutions may also be used and are effective
for inhibition. It will be readily apparent that 100%
sequence identity between the dsRNA and a portion of
the target gene is not absolutely required for
inhibition and the phrase "substantially identical" as
25 used herein is to be interpreted accordingly.
Generally sequences which are substantially identical
will share at least 90%, preferably at least 95% and
more preferably at least 98% nucleic acid sequence
identity. Sequence identity may be conveniently
30 calculated based on an optimal alignment, for example
using the BLAST program accessible at
WWW.ncbi.nlm.nih.gov.

The invention will be further understood with
reference to the following non-limiting Examples,
35 together with the accompanying Figures in which:

Figure 1 is a plasmid map of the vector pGN1

containing opposable T7 promoters flanking a multiple cloning site and an ampicillin resistance marker.

5 Figure 2 is a plasmid map of the vector pGN8 (a genomic fragment of the *C. elegans* *unc-22* gene cloned in pGN1).

10 Figure 3 is a plasmid map of the vector pGN29 containing two T7 promoters and two T7 terminators flanking *Bst*XI sites. This vector permits cloning of DNA fragments linked to *Bst*XI adaptors.

15 Figure 4 is a plasmid map of the vector pGN39 containing two T7 promoters and two T7 terminators flanking attR recombination sites (based on the Gateway™ cloning system of Life Technologies, Inc).

20 Figure 5 is a plasmid map of the vector pGX22 (a fragment of the *C. elegans* gene C04H5.6 cloned in pGN29).

25 Figure 6 is a plasmid map of the vector pGX52 (a fragment of the *C. elegans* gene K11D9.2b cloned in pGN29).

Figure 7 is a plasmid map of the vector pGX104 (a fragment of the *C. elegans* gene Y57G11C.15 cloned in pGN29).

30 Figure 8 is a plasmid map of the vector pGZ8 (a fragment of the *C. elegans* gene T25G3.2 cloned in pGN39).

35 Figure 9 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the

plasmid pGX22.

Figure 10 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX52.

Figure 11 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGXG28.

Figure 12 shows the results of an RNAi experiment in which wild-type (N2) or *nuc-1* strain *C. elegans* in liquid culture were fed with *E. coli* containing the plasmid pGX104

20

Example 1

Influence of genetic background on the efficiency of RNAi in *C. elegans*.

5 Introduction

Various different *C. elegans* strains were fed with different bacteria, to test the possibility of RNAi by feeding *C. elegans* with bacteria that produce dsRNA. The possibility of DNA delivery and dsRNA delivery has
10 previously been envisaged by using different bacterial strains. In this experiment the importance of the *C. elegans* strain as receptor of the dsRNA is also shown.

For this experiment the following *E. coli* strains were
15 used:

1. MC1061: F-araD139 Δ (ara-leu)7696 galE15 galK16
 Δ (lac)X74 rps1 (*Str*^r) hsdR2 (*r*_k⁻ *m*_k⁺) mcrA mcrB1
 - regular host for various plasmids,
 - Wertman et al., (1986) Gene 49:253-262,
 - Raleigh et al., (1989) in Current Protocols in
 Molecular Biology eds. Ausubel et al, Publishing
 associates and Wiley Interscience; New York. Unit
 1.4.
 25
2. B21(DE3): F- ompT(lon) hsdS_B (*r*_B⁻, *m*_B⁻; an *E. coli* B
 strain) with DE3, a λ prophage carrying the T7
 RNA polymerase gene.
 - regular host for IPTG inducible T7 polymerase
 expression,
 - Studier et al. (1990) Meth. Enzymol. 185:60-89
 30
3. HT115 (DE3): F- mcrA mcrb IN(rrnD-rrnE) 1 λ -
 rnc14::tr10 (DE3 lysogen: lacUV5
 promoter-T7polymerase)
 35 - host for IPTG inducible T7 polymerase

expression,

- RNaseIII-,

- Fire A, Carnegie Institution, Baltimore, MD,
Pers. Comm.

5

For this experiment the following *C. elegans* strains
were used:

10

1. *C. elegans* N2: regular WT laboratory strain

15

2. *C. elegans* *nuc-1*(el393): *C. elegans* strain with a
reduced endonuclease activity (>95%); condensed
chromatin persists after programmed cell death;
ingested (bacterial) DNA in the intestinal lumen
is not degraded. Several alleles are described:
el392 (strong allele: has been used for the
experiments described below); n887 (resembles
el392) and n334 (weaker allele)

20

- Stanfield et al. (1998) East Coast Worm
meeting abstract 171,
- Anonymous, Worm Breeder's Gazette 1(1):17b
- Hevelone et al. (1988) Biochem. Genet. 26:447-461
- Ellis et al., Worm breeder's Gazette 7(2):44
- Babu, Worm Breeder's gazette 1(2):10
- 25 - Driscoll, (1996) Brain Pathol. 6:411-425
- Ellis et al., (1991) Genetics 129:79-94

For this experiment the following plasmids were used:

30

pGN1: A vector encoding for ampicillin resistance,
harbouring a multiple cloning site between two
convergent T7 promoters.

35

pGN8: pGN1 containing a genomic fragment of *unc-22*.
Decreased *unc-22* expression via RNAi results in a
"twitching" phenotype in *C. elegans*.

Experimental conditions

12-well micro-titer plates were filled with approximately 2 ml of NGM agar per well (1 litre of NGM agar: 15g Agar, 1g peptone, 3g NaCl, 1ml cholesterol solution (5 mg/ml in EtOH), with sterile addition after autoclaving of 9.5 ml 0.1M CaCl₂, 9.5 ml 0.1 ml MgSO₄, 25 ml 1M KH₂PO₄/K₂HPO₄ buffer pH 6 and 5 ml nystatin solution (dissolved 10 mg/ml in 1:1 EtOH:CH₃COONH₄ 7.5 M).

10

The dried plates were spotted with approximately 50 µl of an overnight culture of bacteria. When IPTG induction was required, 50 µl of a 10 mM stock solution of IPTG was dropped on top of the bacteria lawn, and incubated at 37°C for approximately 4 hours. Individual nematodes at the L4 growth stage were then placed in single wells. In each well 4 nematodes, and the plates were further incubated at 20°C for 6 days to allow offspring to be formed. The F1 offspring of the seeded nematodes were tested for the twitching phenotype.

15

20

Results

Table 1: Percentage of the offspring that show the twitching phenotype

5	MC1061	N2	<i>nuc-1</i>
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	0%	0%
	pGN8 + IPTG	0%	0%
10	BL21 (DE3)		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	20% (+)	>90% (++)
	pGN8 + IPTG	20% (+)	>90% (++±)
15	HT115 (DE3)		
	pGN1	0%	0%
	pGN1 + IPTG	0%	0%
	pGN8	50% (±)	>90% (++)
20	pGN8 + IPTG	80% (++)	>90% (+++)

%: percentage twitchers

+: weak twitching

++: twitching

+++: strong twitching

Conclusions

The experiment with *E. coli* MC1061 shows that no twitching could be observed in this experiment.

30 Neither the N2 nematodes nor the *nuc-1* nematodes showed any twitchers. This is to be expected as *E. coli* MC1061 does not produce any T7 RNA polymerase, and hence the *unc-22* fragment cloned in pGN8 is not

expressed as dsRNA.

5 The experiment with *E. coli* strain BL21(DE3) and
nematode strain N2 shows expected results. BL21(DE3)
harbouring plasmid pGN1 does not result in any
twitching as the pGN1 vector is an empty vector. BL21
(DE3) harbouring PGN8 results in the expression of
unc-22 dsRNA. When this dsRNA is fed to the N2
nematode (indirectly by feeding with the bacteria that
10 produce the dsRNA), this results in a twitching
phenotype, indicating that the dsRNA is able to pass
the gut barrier and is able to perform its interfering
activity.

15 Surprisingly the RNAi effect of the unc-22 dsRNA was
even more pronounced in *C. elegans* strain *nuc-1* than
in the wild type N2 strain. Although one may expect
that the *nuc-1* mutation results in the non-degradation
or at least in a slower degradation of DNA, as the
20 NUC-1 protein is known to be involved in DNase
activity, we clearly observe an enhancement of the
RNAi induced phenotype in *C. elegans* with a *nuc-1*
background. The *nuc-1* mutation has not been cloned
yet, but it has been described that the gene is
25 involved in nuclease activity, and more particularly
DNase activity. If the NUC-1 protein is a nuclease, it
may also have activity on nuclease activity on dsRNA,
which would explain the enhanced RNAi phenotype. The
nuc-1 gene product may be a nuclease, or a regulator
30 of nuclease activity. As the mode of action of RNAi is
still not understood, it is also possible that the
NUC-1 protein is interfering in the mode of action of
RNAi. This would explain why a *nuc-1* mutant is more
sensitive to RNAi.

35 The experiment with the *E. coli* strain HT115 (DE3)

confirms the experiments with the BL21(DE3) strain.
The RNA interference observed with the unc-22 dsRNA is
even higher. In comparison with strain BL21(DE3) this
could be expected, as HT115(DE3) is a RNase III minus
5 strain, and hence is expected to produce larger
amounts of dsRNA, resulting in more prominent RNAi.
This indicates further that the RNAi observed in this
experiment is the result of the dsRNA produced by the
bacteria fed to the *C. elegans*. Feeding *C. elegans*
10 *nuc-1* with HT115(DE3) harbouring pGN8 also results in
higher RNA interference phenotype than feeding the
same bacteria to *C. elegans* wild-type strain N2. Once
again this indicates that improved RNAi can realised
using a nuclease negative *C. elegans* and more
15 particularly with a with the *C. elegans nuc-1* (el392)
strain.

Summary

RNA interference can be achieved in *C. elegans* by
20 feeding the worms with bacteria that produce dsRNA.
The efficiency of this RNA interference is dependent
both on the *E. coli* strain and on the genetic
background of the *C. elegans* strain. The higher the
level of dsRNA production in the *E. coli*, the more
25 RNAi is observed. This can be realised by using
efficient RNA expression systems such as T7 RNA
polymerase and RNase negative strains, such as
RNaseIII minus stains. In this example the level of
dsRNA production varied: HT115(DE3) > BL21(DE3) > MC1061.

30 RNA interference is high in *C. elegans* strains that
are nuclease negative, or that are influenced in their
nuclease activity. This can be realised by using a
mutant strain such as *C. elegans nuc-1*.
35 In this example the sensitivity to RNAi varied:
C. elegans nuc-1 >> *C. elegans* N2

Example 2

Improved RNAi by feeding dsRNA producing bacteria in selected *C. elegans* strains-Comparison of the *nuc-1* strain with several mutants which show improved gut uptake. (designated herein 'gun' mutants). Strains bg77, bg78, bg83, bg84, bg85, bg86, bg87, bg88 and bg89 are typical gun mutant *C. elegans* strains isolated using selection for increased gut uptake (gun phenotype) with the marker dye BCECF-AM.

Experimental conditions:

- 12-well micro-titer plates were filled with approximately 2ml of NGM agar (containing 1ml/l of ampicillin (100µg/ml) and 5 ml of 100mM stock IPTG) per well
- the dried plates were spotted with 25µl of an overnight culture of bacteria (BL21DE3/HT115DE3) containing the plasmids pGN1 (T7prom-T7prom) or pGN8 (T7prom-unc-22-T7prom)
- individual nematodes at the L4 growth stage were then placed in single wells, one nematode per well
- the plates were incubated at 20°C for 6 days to allow offspring to be formed
- the adult F1 offspring of the seeded nematodes were tested for the twitching phenotype

Results:

Table 2:

	20°C/6d	pGN1 HT115DE3	pGN8 BL2DE3	pGN8 HT115DE3	
5	N2	0	1	1	
	bg77	0	1-2	3	
	bg78	0	1	1-2	
	bg83	0	1	1	
10	bg84	0	1-2	3	
	bg85	0	1	3	
	bg86	0	1	2	
	bg87	0	1	1	
	bg88	0	1	1	
15	bg89	0	1	1	

figure legend:

- 0 = no twitching
- 1 = no to weak phenotype
- 2 = clear phenotype
- 3 = strong phenotype

25 Conclusions

- bacterial strain HT115(DE3) shows a better RNAi sensitivity than bacterial strain BL21(DE3)
- the *nuc-1* *C. elegans* strain is a better strain than the Wild-type N2 strain for RNAi sensitivity
- 30 - various gun mutants (improved gut uptake mutants) and more particularly the gun mutant strains bg77, bg84, bg85, bg86 show improved sensitivity to RNAi compared to Wild-type.

A double mutant *C. elegans* strain (nuc-1/gun) shows even greater sensitivity to RNAi compared to wild-type:

- 5 Double mutants were constructed to test the prediction that gun/nuc mutants would even show more enhanced RNAi sensitivity. As an example, the crossing strategy with gun strain bg85 is shown, similar crosses can be conducted with other gun strains, such
10 as bg77, bg84 and bg86.

P0 cross: gun(bg85) x WT males

15 F1 cross: nuc-1 x gun(bg85)/+ males

F2 cross: nuc-1 x gun(bg85)/+; nuc-1/0 males (50%)
nuc-1 x +/+; nuc-1/0 males (50%)

20 F3 single: gun(bg85)/+; nuc-1 hermaphrodites (25%)
+/+; nuc-1 hermaphrodites (75%)

F4 single: gun(bg85); nuc-1 (1/4 of every 4th plate high staining with BCECF)

25 F5 retest: gun(bg85); nuc-1 (100% progeny of F4 singled high staining with BCECF)

To select for the gun phenotype, the fluorescence precursor BCECF-AM is used (obtainable from Molecular probes). The precursor BCECF-AM is cleaved by
30 esterases present in the gut of the worm to generate the dye BCECF which is fluorescent at pH values above 6. This allows selection for worms that have a gun phenotype. BCECF-AM is taken up through the pharynx
35 into the gut lumen and is not fluorescent until it has been cleaved, and the BCECF portion has entered the

cells surrounding the lumen. Wild-type worms will show slower or no increase in BCECF fluorescence.

5 **Example 3**

Improved RNAi feeding in liquid culture using *nuc-1*(el393) *C. elegans*.

Introduction

10 N2 and *nuc-1* *C. elegans* strains were fed with bacteria producing dsRNAs that give lethal phenotypes via RNAi. For this example RNAi was performed in liquid culture instead of on agar plates. We show here for a number of genes that the RNAi effect is more penetrant using
15 the *nuc-1* strain than the N2 strain, and that RNAi can be performed in liquid.

For this experiment the following *E. coli* strains were used:

20

1. HT115 (DE3): F- *mcrA mcrb IN(rrnD-rrnE)* 1 λ -
 rncl4::trl0 (DE3 lysogen: lacUV5 promoter -T7
 polymerase)
 - host for IPTG inducible T7 polymerase expression
25 - RNaseIII⁻
 - Fire A, Carnegie Institution, Baltimore, MD,
 Pers. Comm.

25

For this experiment, following *C. elegans* strains were
30 used:

30

1. *C. elegans* N2: regular WT laboratory strain
2. *C. elegans nuc-1*(el393): *C. elegans* strain with a
35 reduced endonuclease activity (>95%); condensed
 chromatin persists after programmed cell death;

35

ingested (bacterial) DNA in the intestinal lumen is not degraded. Several alleles are described:

e1392 (strong allele: has been used for the experiments described below); n887 (resembles e1392) and n334 (weaker allele)

5

- Stanfield et al. (1998) East Coast Worm meeting abstract 171

- Anonymous, Worm Breeder's Gazette 1(1):17b

- Hevelone et al. (1988) Biochem. Genet. 26:447-461

10

- Ellis et al., Worm breeder's Gazette 7(2):44

- Babu, Worm Breeder's gazette 1(2):10

- Driscoll, (1996) Brain Pathol. 6:411-425

- Ellis et al., (1991) Genetics 129:79-94

15

For this experiment, the following plasmids that all give lethal phenotypes in *C. elegans* via RNAi were used:

20

pGX22: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid C04H5.6 corresponding to a member of the RNA helicase family.

25

pGX52: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid K11D9.2b corresponding to sarco/endoplasmic Ca²⁺ ATPase also known as SERCA.

30

pGZ18: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid T25G3.2 corresponding to a chitin like synthase gene.

35

pGX104: a vector encoding ampicillin resistance, containing a genomic fragment of cosmid Y57G11C.15 corresponding to sec-61, a transport protein.

Experimental conditions

- 5 - 1 ml overnight cultures of HT115 (DE3) bacteria containing the plasmids pGX22, pGX52, pGZ18 or pGX104 respectively were pelleted and resuspended in S-complete medium, containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 10 - 10 µl of this bacterial solution was transferred to a 96-well microtiter plate already filled with 100 µl S-complete containing 1ml/l of ampicillin (100 µg/ml) and 1ml/l of 1000mM IPTG.
- 15 - 3 nematodes at the L1 growth stage of N2 and nuc-1 strain were then placed in single wells, 3 L1's per well. Per experimental set up, 16 wells were used (n=16).
- 20 - the plates were incubated at 25°C for 5 days to allow offspring to be formed.
- 25 - the plates were visually checked and the following phenotypes could be scored per individual well:
 - 25 no effect: L1's developed to adults and gave normal offspring.
 - 30 no F1 offspring: L1's developed to adults and gave no offspring.
 - 30 acute lethal: original L1 did not mature and died.

Results

- 35 The results of this experiment are illustrated graphically in Figures 9 to 12. Data are expressed as

a percentage of the total (n=16) on the y-axis for both N2 and nuc-1 strains.

Conclusions

5 The following genes were tested in this liquid RNAi assay:

- C04H5.6: an RNA helicase. RNAi of this gene interferes with the generation of offspring.
- 10 - SERCA: a sarco/endoplasmic Ca^{2+} ATPase. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- T25G3.2: a chitin like synthase gene. RNAi of
15 this gene causes dead eggs.
- sec-61: a transport protein. A strong RNAi phenotype causes an acute lethal phenotype. A less penetrant RNAi effect results in loss of offspring.
- 20 - RNAi can be performed under liquid conditions.

As in the previous examples this set of experiments shows that the nuc-1 *C. elegans* strain is more sensitive to RNAi than the wild-type N2 strain. This
25 is most clear for less penetrant phenotypes such as SERCA and chitin synthase. For strong RNAi phenotypes like the helicase and Sec-61 the difference between the N2 wild-type strain and the nuc-1 stain is less pronounced.

30

Example 4

Cloning of pGX22, pGX52, pGZ18 and pGX104 for RNAi

A set of primers for each gene was designed on the basis of sequence data available in the publicly accessible *C. elegans* sequence database (Acedb).

The cosmid names relate to:

1. C04H5.6=member of RNA helicase
2. K11D9.2b=SERCA
3. Y57G11C.15=transport protein sec-61
4. T25G3.2=chitin synthase like

The following primer sequences were designed:

1. C04H5.6F 5'-TGCTCAGAGAGTTTCTCAACGAACC-3'
C04H5.6R 5'-CAATGTTAGTTGCTAGGACCACCTG-3'
2. K11D9.2bF 5'-CAGCCGATCTCCGTCTTGTG-3'
K11D9.2bR 5'-CCGAGGGCAAGACAACGAAG-3'
3. Y57G11C.15F 5'-ACCGTGGTACTCTTATGGAGCTCG-3'
Y57G11C.15R 5'-TGCAGTGGATTGGGTCTTCG-3'
4. T25G3.2F
5'-GGGGACAAGTTTGTACAAAAAGCAGGCTATGCCAAGTACATGTCGATTGCG-3'

T25G3.2R
5'-GGGGACCACTTTGTACAAGAAAGCTGGGTGGAGAAGCATTCCGAGAGTTTG-3'

PCR was performed on genomic DNA of N2 strain *C. elegans* to give PCR products of the following sizes:

- 1326bp for C04H5.6
1213bp for K11D9.2b

1024bp for Y57G11C.15

1115bp for T25G3.2

- 5 The PCR fragments of C04H5.6, K11D9.2b and Y57G11C.15 were linked to *Bst*XI adaptors (Invitrogen) and then cloned into the pGN29 vector cut with *Bst*XI. pGN29 contains two T7 promoters and two T7 terminators flanking a cloning site which is adapted for facilitated cloning of PCR fragments, comprising a
- 10 stuffer DNA flanked by two *Bst*XI sites (see schematic Figure 3). The resulting plasmids were designated pGX22 (C04H5.6), pGX52 (K11D9.2b) and pGX104 (Y57G11C.15).
- 15 The PCR fragment of T25G3.2 was cloned into pGN39 via recombination sites based on the GATEWAY™ cloning system (Life Technologies, Inc). pGN39 contains two T7 promoters and two T7 terminators flanking a cloning site which facilitates "High Throughput" cloning based
- 20 on homologous recombination rather than restriction enzyme digestion and ligation. As shown schematically in Figure 4, the cloning site comprises *attR1* and *attR2* recombination sites from bacteriophage lambda flanking a gene which is lethal to *E. coli*, in this
- 25 case the *ccdB* gene. This cloning site is derived from the Gateway™ cloning system commercially available from Life Technologies, Inc. The Gateway™ cloning system has been extensively described by Hartley et al. in WO 96/40724 (PCT/US96/10082).
- 30

Example 5

Selecting *C. elegans* mutations for increased gut uptake (gun) using marker dye BCECF-AM and *unc-31* as background.

5

The screen was performed in *unc-31(e928)* mutant background, to ensure high amounts of dye in the gut lumen, since *unc-31* mutations show constitutive pharyngeal pumping. The dye (BCECF-AM: 2',7' bis (2-carboxyethyl)-5-(and-6)-carboxyfluorescein, acetoxymethylester); obtained from Molecular Probes, is cleaved by intracellular esterases. Fluorescence accumulates in the gut cells upon passage through the apical gut membrane.

10
15**Mutagenesis**

Day 1: *unc-31* L4 staged worms were mutagenised with EMS (final concentration 50 mM) for 4 hours

Day 2: P0 was divided over several large agar plates

20

Day 6: F1's were collected and dropped on large plates. The number of eggs the F1's layed were checked every hour and the F1's were removed when 10-20 eggs per F1 were counted

25

Day 10: F2 adults were collected and screened with BCECF-AM. Mutations with increased staining of the gut cells after 15-30 minutes exposure to the dye were selected and singled on small agar plates.

30

About 50 initial positives gave progeny which was retested with BCECF-AM (2x) and leucine CMB (1x) 9 of the 50 strains were kept (2 strains : 3 times positive, 7 other strains : twice positive)

35

Table 3: Isolation of mutations for increased staining with BCECF-AM

	Total P0	Total F1	Total F2	screened chromosomes	number of strains isolated
5	(counted)	(estimated)	(calculated)	(estimated)	(counted)
	2251	55618	222472	100000	9

Outcrossing, backcrossing and double construction

- 10 1. backcrossing *unc-31; gun* --> *unc-31; gun*
 - *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains segregating 1/4 *unc*
- 15 - stain *unc* strains with BCECF-AM
 - from positive strains pick *unc* homozygous
 - retest 100 % *unc* strains with BCECF-AM
 - kept 1 strain (backcrossed)
- 20 2. *unc-31* background was crossed out-->+; *gun*
 - *unc-31; gun* x WT males
 - singled 2x5 WT hermaphrodites F1s (*unc-31/+; gun/+*)
 - singled 50 WT hermaphrodites F2s (1/4 homozygous)
 - select strains which did not segregate *unc* F3s
- 25 anymore
 - stain non *unc* strains with BCECF-AM
 - 7 positive strains were retested with BCECF-AM and finally 1 was selected and kept (outcrossed)
- 30 3. +; *gun* (1x outcrossed) were 2 times backcrossed-->+; *gun* (3x backcrossed)
 - *gun* x WT males
 - WT hermaphrodites x F1 males (*gun/+*)
 - singled 10 WT hermaphrodites F2s (1/2 heterozygous)
- 35 - singled 50 WT hermaphrodites F3s (1/8 homozygous)

- 31 -

- stain strains with BCECF-AM- retested positives with BCECF-AM and finally 1 was selected and kept

4. *gun* (3x backcrossed) were crossed with *nuc-1(X)*
 5 mutant--> *gun; nuc-1*
 - *gun* x WT males
 - *nuc-1* x *gun/+* males
 - *nuc-1* x *gun/+; nuc-1/0* or *+/+; nuc-1/0* males
 - singled 10 WT hermaphrodite progeny (*nuc-1*
 10 homozygous, $\frac{1}{2}$ heterozygous *gun*)
 - singled 40 WT hermaphrodite progeny (1/8 homozygous *gun*)
 - stain strains with BCECF-AM
 - retested positives with BCECF-AM and finally 1 was
 15 selected and kept

Table 6: Strains derived from *gun* mutations

20	gun	unc-31; gun		unc-31; gun		+; gun			gun; nuc-1
		original isolate		backcrossed (1x)		outcrossed (1x)		3x b.c.	from 3x b.c.
	allele number	isolation number	strain number	isolation number	strain number	isolation number	strain number	strain number	strain number
	bg77	31.4	UG 510	31.4.46.1	UG 556	31.4.34	UG 563	UG 674	UG 777
25	bg78	37.5	UG 511	37.5.46.4	UG 557	37.5.15	UG 564	UG 675	—
	bg83	10.2	UG 543	10.2.11	UG 600	10.2.21	UG 586	UG 676	—
	bg84	7.2	UG 544	7.2.10	UG 601	7.2.15	UG 589	UG 677	UG 774
	bg85	11.5	UG 545	11.5.29.2	UG 602	2x b.c.	UG 717		UG 775
	bg86	42.1	UG 546	42.1.4.5	UG 603	42.1.18	UG 587	UG 678	UG 776
30	bg87	7.1	UG 547	7.1.8.3	UG 604	7.1.22	UG 585	UG 679	—
	bg88	5.3	UG 548	5.3.9	UG 605	5.3.18	UG 584	UG 680	—
	bg89	23.4	UG 549	23.4.13.5	UG 606	23.4.3	UG 588	UG 671	—

SEQUENCE LISTING:

SEQ ID NO: 1 complete sequence of pGN1

5 SEQ ID NO: 2 complete sequence of pGN8

SEQ ID NO: 3 complete sequence of pGN29

SEQ ID NO: 4 complete sequence of pGN39

10 SEQ ID NO: 5 complete sequence of pGX22

SEQ ID NO: 6 complete sequence of pGX52

15 SEQ ID NO: 7 complete sequence of pGX104

SEQ ID NO: 8 complete sequence of pGZ8

SEQ ID NO: 9 primer C04H5.6F

20 SEQ ID NO: 10 primer C04H5.6R

SEQ ID NO: 11 primer K11D9.2bF

25 SEQ ID NO: 12 primer K11D9.2bR

SEQ ID NO: 13 primer Y57G11C.15F

SEQ ID NO: 14 primer Y57G11C.15R

30 SEQ ID NO: 15 primer T25G3.2F

SEQ ID NO: 16 primer T25G3.2R

Claims:

1. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to
5 said nematode worm a food organism which is capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene following ingestion of the food organism by the nematode, wherein the nematode
10 has a non wild-type genetic background selected to provide increased sensitivity to RNA interference as compared to wild type.
2. A method as claimed in claim 1 wherein the
15 nematode is a microscopic nematode.
3. A method as claimed in claim 2 wherein the nematode is from the genus *Caenorhabditis*.
- 20 4. A method as claimed in claim 3 wherein the nematode is *C. elegans*.
5. A method as claimed in any one of claims 1 to 4 wherein the nematode has a mutant genetic
25 background.
6. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits reduced activity of one or more nucleases compared to wild
30 type.
7. A method as claimed in claim 6 wherein the nematode is *C. elegans* strain *nuc-1*.
- 35 8. A method as claimed in claim 5 wherein the nematode is a mutant strain which exhibits increased

gut uptake compared to wild type.

9. A method as claimed in claim 8 wherein the nematode is mutant *C. elegans* strain bg85.

5

10. A method as claimed in claim 5 wherein the nematode is a mutant strain having at least one mutation which results in reduced nuclease activity compared to wild type and at least one mutation which results in increased gut uptake compared to wild type.

10

11. A method as claimed in claim 10 wherein the nematode is a mutant *C. elegans* strain having the *nuc-1* mutation and the bg85 mutation.

15

12. A method as claimed in any one of the preceding claims wherein the food organism has been engineered to express a double-stranded RNA.

20

13. A method as claimed in any one of the preceding claims wherein the food organism is a bacterium.

14. A method as claimed in claim 13 wherein the food organism is *E. coli*.

25

15. A method as claimed in any one of the preceding claims wherein the food organism has been genetically modified to express a double-stranded RNA having a nucleotide sequence substantially identical to a portion of said target gene.

30

16. A method as claimed in claim 15 wherein the food organism contains a DNA vector, the vector comprising a promoter or promoters orientated relative to a DNA sequence such that they are capable of

35

initiating transcription of said DNA sequence to RNA capable of forming a double-stranded structure upon binding of an appropriate RNA polymerase to said promoter or promoters.

5

17. A method as claimed in claim 25 wherein the vector comprises two promoters flanking the DNA sequence.

10

18. A method as claimed in claim 26 wherein the two promoters are identical.

15

19. A method as claimed in claim 25 wherein the vector comprises a single promoter and further comprises said DNA sequence in a sense and an antisense orientation relative to said promoter.

20

20. A method as claimed in any one of claims 16 to 20 wherein the nematode or the food organism is adapted to express an RNA polymerase capable of initiating transcription from said promoter or promoters.

25

21. A method as claimed in any one of claims 16 to 20 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

30

22. A method as claimed in any one of claims 1 to 21 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

35

23. A method of inhibiting expression of a target gene in a nematode worm comprising feeding to said nematode worm a food organism capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a

portion of said target gene following ingestion of the food organism by the nematode, wherein the food organism carries a modification selected to provide increased expression or persistence of the doubled-stranded RNA compared to a food organism which does not carry the modification.

24. A method as claimed in claim 23 wherein the food organism is a bacterium.

25. A method as claimed in claim 24 wherein the bacterium is an *E. coli* strain.

26. A method as claimed in claim 25 wherein the *E. coli* strain is an RNase III minus strain or any other RNase negative strain.

27. A method as claimed in any one of claims 23 to 26 wherein the step of feeding said food organism to said nematode worm is carried out in liquid culture.

28. A method of inhibiting expression of a target gene in a nematode worm comprising introduction of a DNA capable of producing a double-stranded RNA structure having a nucleotide sequence substantially identical to a portion of said target gene in said nematode, wherein the nematode is one which exhibits increased gut uptake compared to wild type.

29. A method as claimed in claim 28 wherein the nematode is a microscopic nematode.

30. A method as claimed in claim 29 wherein the nematode is from the genus *Caenorhabditis*.

31. A method as claimed in claim 30 wherein the

nematode is *C. elegans*.

32. A method as claimed in any one of claims 28
to 31 wherein the nematode has a mutant genetic
5 background.

33. A method as claimed in claim 32 wherein the
nematode is mutant *C. elegans* strain bg85.

10 34. A method as claimed in any one of claims 28
to 33 wherein the DNA capable of producing a double-
stranded RNA structure is a vector comprising a
promoter or promoters orientated relative to a DNA
sequence such that they are capable of initiating
15 transcription of said DNA sequence to RNA capable of
forming a double-stranded structure upon binding of an
appropriate RNA polymerase to said promoter or
promoters.

20 35. A method as claimed in claim 34 wherein the
vector comprises two promoters flanking the DNA
sequence.

25 36. A method as claimed in claim 35 wherein the
two promoters are identical.

30 37. A method as claimed in claim 34 wherein the
vector comprises a single promoter and further
comprises said DNA sequence in a sense and an
antisense orientation relative to said promoter.

35 38. A method as claimed in any one of claims 34
to 37 wherein the nematode is adapted to express an
RNA polymerase capable of initiating transcription
from said promoter or promoters.

39. A method as claimed in any one of claims 34

to 38 wherein the RNA polymerase is T7, T3 or SP6 polymerase.

FIG. 1.




FIG. 2.




FIG. 3.




FIG. 4.




FIG. 5.




FIG. 6.




FIG. 7.




FIG. 8.




FIG. 9.




FIG. 10.




FIG. 11.





FIG. 12.



1
SEQUENCE LISTING

<110> DEVGEN NV

<120> IMPROVEMENTS RELATING TO DOUBLE-STRANDED RNA INHIBITION

<130> SCB/53711/001

<140>

<141>

<160> 14

<170> PatentIn Ver. 2.0

<210> 1

<211> 3216

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN1

<400> 1

```
gagtgcacca tatgcggtgt gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgttaa tattttgtta aaattcgcgt taaatatttg ttaaatacagc 120
tcatttttta accaataggc cgaaatcggc aaaatccctt ataaatcaaa agaatagacc 180
gagatagggt tgagtgttgt tccagtttgg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aagggcgaaa aaccgtctat cagggcgatg gccactacg tgaaccatca 300
cccaaataca gttttttgcg gtcgaggtgc cgtaaagctc taaatcggaa ccctaaaggg 360
agcccccgat ttagagcttg acgggggaaag ccggcgaaag tggcgagaaa ggaaggggaag 420
aaagcgaaaag gagcggggcgc tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacacccg ccgcgcttaa tgcgccgcta cagggcgctg ccattcgcca ttcaggctgc 540
gcaactgttg ggaagggcga tcggtgcggg cctcttcgct attacgccag ctggcgaaaag 600
ggggatgtgc tgcaaggcga ttaagttggg taacgccagg gttttcccag tcacgacgtt 660
gtaaaacgac ggccagtgaa ttgtaatacg actcactata gggcgaaattc gagctcggta 720
cccggggatc ctctagagtc gaaagcttct cgccctatag tgagtcgtat tacagcttga 780
gtattctata gtgtcaccta aatagcttgg cgtaatcatg gtcatagctg tttcctgtgt 840
gaaattgtta tccgctcaca attccacaca acatacgagc cggaagcata aagtgtaaag 900
cctgggggtgc ctaatgagtg agctaactca cattaattgc gttgcgctca ctgcccgtt 960
tccagtcggg aaacctgtcg tgccagctgc attaatgaat cggccaacgc gcggggagag 1020
gcggtttgcg tattgggcgc tcttcgctt cctcgctcac tgactcgctg cgctcggctg 1080
ttcggtgcg gcgagcggtg tcagctcact caaaggcggg aatacggtta tccacagaat 1140
caggggataa cgcaggaaaag aacatgtgag caaaaggcca gcaaaaggcc aggaaccgta 1200
aaaaggccgc gttgctggcg tttttcgata ggctccgccc ccctgacgag catcacaaaa 1260
atcgacgctc aagtcagagg tggcgaaacc cgacaggact ataaagatac caggcgtttc 1320
cccctggaag ctccctcgtg cgctctcctg ttccgaccct gccgttacc ggatacctgt 1380
ccgcctttct cccttcggga agcgtggcgc tttctcatag ctacgctgt aggtatctca 1440
gttcggtgta ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagccc 1500
accgctgcgc cttatccggt aactatcgtc ttgagtccaa cccgtaaga cacgacttat 1560
cgccactggc agcagccact ggtaacagga ttagcagagc gaggtatgta ggcggtgcta 1620
cagagttctt gaagtgttg cctaactacg gctacactag aaggacagta tttggtatct 1680
gcgctctgct gaagccagtt accttcggaa aaagagttgg tagctcttga tccggcaaac 1740
aaaccaccgc tggtagcggg ggtttttttg tttgcaagca gcagattacg cgcagaaaaa 1800
aaggatctca agaagatcct ttgatctttt ctacgggggtc tgacgctcag tggaacgaaa 1860
actcacgtta agggattttg gtcagtgcgt tatcaaaaag gatcttcacc tagatccttt 1920
taaattaaaa atgaagtttt aaatcaatct aaagtatata tgagtaaact tggctctgaca 1980
```

2

gttaccaatg	cttaatcagt	gaggcaccta	tctcagcgat	ctgtctatatt	cgttcatcca	2040
tagttgctg	actccccgtc	gtgtagataa	ctacgatacg	ggagggctta	ccatctggcc	2100
ccagtgcctg	aatgataccg	cgagacccac	gctcaccggc	tccagattta	tcagcaataa	2160
accagccagc	cggaagggcc	gagcgcagaa	gtggtcctgc	aactttatcc	gcctccatcc	2220
agtctattaa	ttgttgccgg	gaagctagag	taagtagttc	gccagttaat	agtttgcgca	2280
acgttggttg	cattgctaca	ggcatcgtgg	tgtcacgctc	gtcgtttggt	atggcttcat	2340
tcagctccgg	ttcccaacga	tcaaggcgag	ttacatgata	ccccatgttg	tgcaaaaaag	2400
cggttagctc	cttcgggtcct	ccgatcgttg	tcagaagtaa	gttgcccgca	gtgttatcac	2460
tcattggttat	ggcagcactg	cataattctc	ttactgtcat	gccatccgta	agatgctttt	2520
ctgtgactgg	tgagtactca	accaagtcac	tctgagaata	ccgcgcccgg	cgaccgagtt	2580
gctcttgccc	ggcgtcaata	cgggataata	gtgtatgaca	tagcagaact	ttaaaagtgc	2640
tcattcattg	aaaacgttct	tcggggcgaa	aactctcaag	gatcttaccg	ctgttgagat	2700
ccagttcgat	gtaaccact	cgtgcaccca	actgatcttc	agcatctttt	actttcacca	2760
gcgtttctgg	gtgagcaaaa	acaggaaggc	aaaatgccgc	aaaaaaggga	ataagggcga	2820
cacggaaatg	ttgaatactc	atactcttcc	tttttcaata	ttattgaagc	atttatcagg	2880
gttattgtct	catgagcgga	tacatatattg	aatgtattta	gaaaaataaa	caaatagggg	2940
ttccgcgcac	atttccccga	aaagtgccac	ctgacgtcta	agaaaccatt	attatcatga	3000
cattaaccta	taaaaatagg	cgtatcacga	ggccctttcg	tctcgcgcgt	ttcgggtgatg	3060
acggtgaaaa	cctctgacac	atgcagctcc	cggagacggg	cacagcttgt	ctgtaagcgg	3120
atgccgggag	cagacaagcc	cgtcagggcg	cgtcagcggg	tgttggcggg	tgtcggggct	3180
ggcttaacta	tgccgcatca	gagcagattg	tactga			3216

<210> 2

<211> 4620

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGN8

<400> 2

gatccgaatc	tccatgtctg	ttaacagcct	tgacacggaa	tttatattca	tgcccttgag	60
tcaaactcgtc	aacgtggaag	ttggatcctt	tgctctctcc	gcaagcagtc	catctgccag	120
tggcagcacc	ttgcttttca	atgacatagt	gactgatttc	agctcctcca	tcattctctg	180
gttccttcca	tgcaagatca	catccatcct	tgacaatatt	agtgcacatc	agaggccac	240
gtgggcttga	tgatgatca	agaacagtaa	ccttcacttc	agcagtgtca	gttccattct	300
cgttctctgc	cttgatgata	taggttcctg	tatccgaacg	caaagctctc	ttcacatgga	360
atttagtctt	gccgtcttca	ttgttcaact	tcatacgatc	atcagattcg	actggtgttc	420
cttcgaaagt	ccaagtaatt	gttggaagttg	gttcaccact	gactggaatg	ttcaatgaga	480
agtcttgctc	agccttgacc	ttgatttctt	gaatcgagtt	acgatcgatg	actggtggaa	540
ctataattta	attcaatgat	tattagtaat	tgatttagac	tcttaccatt	tctagccttt	600
gcaacagctg	atgctgaatc	agatggatct	cccaatcctg	ccttggttctt	ggcacggatt	660
ctgaattcgt	actttgatcc	ttccttgaga	tttccaacag	tagcattcgt	ttgtccagct	720
ggaacatgag	caacgtcatt	ccagaatggc	gagaactcgt	ccttcactctc	aacaacgtat	780
tcctcgattg	gagcaccacc	gtcgtttgct	ggtggcttcc	attcaaggtc	aacatgatcc	840
ttatcccaat	cagtaatttc	aggagcattt	gtcttctctg	gcttggtcaa	tgatctttg	900
gcaagtgtgg	ttccgaagg	ctccaatgga	tcggactctc	cttcagcatt	gacggcagcg	960
acacggaact	gaaaatcaaa	atggtttagg	caattgagtt	caagattaaa	aaattctcac	1020
tttatattca	tgtccaggaa	taagaccgtc	aacaacagct	gtagtcttat	ctccagcgac	1080
ctttgcagct	ggaacccatc	ttccacttgc	agtatcgatc	ttttcgatca	catagttttc	1140
aattggaata	cctccatcat	catctggtgc	acgccaattc	aaagtgcac	gatcaccatg	1200
aacatcgga	acatctaatt	gaccatttgg	agaagttggc	ttgtctgaaa	atttaaaata	1260
taaccaaat	aattgaagaa	aactaatgct	caccaataac	attgatctta	acagttgctt	1320
catcttctcc	atttgcatgg	acagctttga	tagtgaaagt	tccactgtct	ccacgttcca	1380
tttgcttca	aaccagcttt	gattggtatt	ctgggttatc	aagcttctcg	ccctatagtg	1440
agtcgtatta	cagcttgagt	attctatagt	gtcacctaaa	tagcttggcg	taatcatggt	1500
catagctgtt	tcctgtgtga	aattgttatc	cgctcacaat	tccacacaac	atacgagccg	1560
gaagcataaa	gtgtaaagcc	tggggtgcct	aatgagttag	ctaactcaca	ttaattgcgt	1620

tgcgctcact gcccgctttc cagtcgggaa acctgtcgtg ccagctgcat taatgaatcg 1680
gccaacgcgc ggggagagggc ggtttgcgta ttgggcgctc ttccgcttcc tcgctcactg 1740
actcgctgcg ctccggtcgtt cggctgcggc gagcgggtatc agctcactca aaggcggtaa 1800
tacggttatc cacagaatca ggggataacg caggaaagaa catgtgagca aaaggccagc 1860
aaaaggccag gaaccgtaaa aaggccgcgt tgctggcggtt ttccgatagg ctccgcccc 1920
ctgacgagca tcacaaaaat cgacgctcaa gtcagaggtg gcgaaacccg acaggactat 1980
aaagatacca ggcgtttccc cctggaagct ccctcgtgcg ctctcctggt cccaccctgc 2040
cgcttaccgg atacctgtcc gcctttctcc ctccgggaag cgtggcgctt tctcatagct 2100
cacgctgtag gtatctcagt tcggtgtagg tcgttcgctc caagctgggc tgtgtgcacg 2160
aacccccgt tcagcccgac cgctgcgcct tatccggtaa ctatcgtctt gagtccaacc 2220
cggtaagaca cgacttatcg ccactggcag cagccactgg taacaggatt agcagagcga 2280
ggtatgtagg cgggtgctaca gagttcttga agtgggtggc taactacggc tacactagaa 2340
ggacagtatt tggatctgc gctctgctga agccagttac ctccgaaaaa agagttggta 2400
gctcttgatc cggcaacaa accaccgctg gtagcgggtg tttttttgtt tgcaagcagc 2460
agattacgcg cagaaaaaaa ggatctcaag aagatccttt gatcttttct acggggtctg 2520
acgctcagtg gaacgaaaac tcacgttaag ggatttttgt catgagatta tcaaaaagga 2580
tcttcacctg gatcctttta aattaaaaat gaagttttta atcaatctaa agtatatatg 2640
agtaaaactg gtctgacagt taccaatgct taatcagtga ggcacctatc tcagcgatct 2700
gtctatttctg ttcattccata gttgcctgac tccccgctcg gtagataact acgatacggg 2760
agggcttacc atctggcccc agtgctgcaa tgataccgcg agaccacgc tcaccggctc 2820
cagatttatc agcaataaac cagccagccg gaagggccga gcgcagaagt ggtcctgcaa 2880
ctttatccgc ctccatccag tctattaatt gttgccggga agctagagta agtagttcgc 2940
cagttaatag tttgcgcaac gttgttgcca ttgctacagg catcgtggtg tcacgctcgt 3000
cgtttggtat ggcttcattc agctccggtt cccaacgatc aaggcgagtt acatgatccc 3060
ccatgttgtg caaaaaagcg gttagctcct tcggtcctcc gatcgttgtc agaagtaagt 3120
tggccgcagt gttatcactc atgggttatgg cagcactgca taattctctt actgtcatgc 3180
catccgtaag atgcttttct gtgactggtg agtactcaac caagtcattc tgagaatacc 3240
gcgcccggcg accgagttgc tcttgcccgg cgtcaatacg ggataatagt gtatgacata 3300
gcagaacttt aaaagtgtc atcattggaa aacgttcttc ggggcgaaaa ctctcaagga 3360
tcttacgct gttgagatcc agttcgatgt aaccactcgc tgcacccaac tgatcttcag 3420
catcttttac tttcaccagc gtttctgggt gagcaaaaac aggaaggcaa aatgccgcaa 3480
aaaagggaat aaggcgacga cggaaatgtt gaatactcat actcttcctt tttcaatatt 3540
attgaagcat ttatcagggc tattgtctca tgagcggata catatttgaa tgtatttaga 3600
aaaataaaca aataggggtt ccgcgcacat ttccccgaaa agtgccacct gacgtctaag 3660
aaaccattat tatcatgaca ttaacctata aaaataggcg tatcacgagg ccctttcgtc 3720
tcgcgcgttt cgggtgatgac ggtgaaaacc tctgacacat gcagctcccg gagacggtca 3780
cagcttgtct gtaagcggat gccgggagca gacaagcccg tcagggcgcg tcagcgggtg 3840
ttggcgggtg tcggggctgg cttaactatg cggcatcaga gcagattgta ctgagagtgc 3900
accatatgcg gtgtgaaata ccgcacagat gcgtaaggag aaaataccgc atcaggcgaa 3960
attgtaaacg ttaatatatt gttaaaattc gcgttaaata tttgttaaata cagctcattt 4020
tttaaccaat aggccgaaat cggcaaaatc cttataaat caaaagaata gaccgagata 4080
gggttgagtg ttgttccagt ttggaacaag agtccactat taaagaacgt ggactccaac 4140
gtcaaagggc gaaaaaccgt ctatcagggc gatggcccac tacgtgaacc atcacccaaa 4200
tcaagttttt tgcggtcgag gtgccgtaaa gctctaaatc ggaaccctaa agggagcccc 4260
cgatttagag cttgacgggg aaagccggcg aacgtggcga gaaaggaagg gaagaaagcg 4320
aaaggagcgg gcgctagggc gctggcaagt gtagcggtea cgctgcgcgt aaccaccaca 4380
cccgcgcgc ttaatgcgcc gctacagggc gcgtccattc gccattcagg ctgcgcaact 4440
gttggaagg gcgatcgggt cgggcctctt cgctattacg ccagctggcg aaagggggat 4500
gtgctgcaag gcgattaagt tgggtaacgc cagggttttc ccagtcacga cgttgtaaaa 4560
cgacggccag tgaattgtaa tacgactcac tataggcgca attcgagctc ggtaccggg 4620

<210> 3

<211> 4756

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX22

<400> 3

tgctcagaga	gtttctcaac	gaacccgatt	tggttagtta	taggtaattt	ttagaacatt	60
tacaaaaaca	gcaaaaaaac	caaacattca	ggatttttgt	ttttaattaa	gaaaaaaatc	120
gatcgctctt	aaattttaat	caatacttcg	aataaaccca	aaaaaaaacg	aaaaaaaatc	180
ctgtttccag	tgtaatgatg	attgacgagg	ctcacgaacg	tactctacac	acggatatte	240
tattcgggtt	agtcaaagat	attgcaagat	tccgaaagga	tttgaagctt	ctcatctctt	300
ctgcaacact	tgacgctgaa	aagttctcca	gtttcttcga	cgacgctccg	attttccgaa	360
ttccgggacg	cagattcccg	gtggacattt	actatacaca	ggctcccga	gcggactacg	420
tcgacgcggc	tatcgtcaca	attatgcaga	ttcacttgac	ccagccactt	cccggcgata	480
ttttggtatt	tctgactggt	caggaagaaa	tcgaaactgt	acaggaagca	cttatggaa	540
ggtcgaaagc	actgggatcg	aagattaagg	agcttattcc	gctgccgggt	tatgcgaatt	600
tgcccagtg	tttgcaggcg	aagattttcg	agccaacgcc	gaaagatg	agaaaggtag	660
atttttctta	caaatttttt	ccaaaaaaa	atccgagaaa	aatctacaaa	atttcaggca	720
aaaactgttt	cattttattc	ctaactagtt	ttttagcaaa	cgttttagatt	taacaaaact	780
gaacaaattt	gaagttttcc	aatttaaaaa	ataaatgttt	cggaaagt	attgaaaaat	840
ctgaaattgc	tatcctctcg	tatctgcaaa	aaaaacactt	taaaaaatgc	tctgttcttt	900
gaaaatttct	aaactgaaaa	atgtgaaatt	tctgaaatt	gtgataattt	tataaaattt	960
tatagaaaat	gtaagcattc	cagaaaaata	tcaaaaattt	cgagaaaatt	ctgaaaaaat	1020
ccagaaatat	taacagaaaa	aaaatctttt	gaaacatctg	aaaattaaaa	taaattgaat	1080
ttacattttt	ttttttggga	tttcttaaaa	atcactatga	atttaccact	aaattttttg	1140
caaaaaatta	tttttttaat	ttcaaagaaa	aagcaaagaa	ttttaaaata	tcaaaaagtc	1200
caaatttggt	tcggtgaatt	tttaaaataa	cattttcaag	ataattttta	gttaatcaaa	1260
acattccacg	cattttctagt	ttcccaaatt	tctctaaatt	tcaggtggtc	ctagcaacta	1320
acattgccag	cacaatggat	ctcgagggat	cttccatacc	taccagttct	gcgcctgcag	1380
gtcgcggccg	cgactctcta	gacgcgtaag	cttactagca	taaccctctg	gggcctctaa	1440
acgggtcttg	aggggttttt	tgagcttctc	gccctatagt	gagtcgtatt	acagcttgag	1500
tattctatag	tgtcacctaa	atagcttggc	gtaatcatgg	tcatactgtg	ttcctgtgtg	1560
aaattgttat	ccgctcacia	ttccacacia	catacgagcc	ggaagcataa	agtgtaaagc	1620
ctggggtgcc	taatgagtga	gctaactcac	attaattgcg	ttgcgctcac	tgcccgtttt	1680
ccagtcggga	aacctgtcgt	gccagctgca	ttaatgaatc	ggccaacgcg	cggggagagg	1740
cggtttgctg	attgggcgct	cttccgcttc	ctcgctcact	gactcgctgc	gctcggctcg	1800
tcggctgcgg	cgagcgggat	cagctcactc	aaaggcggta	atacggttat	ccacagaatc	1860
aggggataac	gcaggaaaga	acatgtgagc	aaaaggccag	caaaaggcca	ggaaccgtaa	1920
aaaggccgcg	ttgctggcgt	ttttcgatag	gctccgcccc	cctgacgagc	atcacaaaaa	1980
tcgacgctca	agtcagaggt	ggcgaaaccc	gacaggacta	taaagatacc	aggcgtttcc	2040
ccctggaagc	tccctcgtgc	gctctcctgt	tcgaccctg	ccgcttaccg	gatacctgtc	2100
cgcctttctc	ccttcgggaa	gcgtggcgct	ttctcatagc	tcacgctgta	ggtatctcag	2160
ttcgggtgtg	gtcgttcgct	ccaagctggg	ctgtgtgcac	gaaccccccg	ttcagcccga	2220
ccgctgcgcc	ttatccggtg	actatcgtct	tgagtccaac	ccggttaagac	acgacttate	2280
gccactggca	gcagccactg	gtaacaggat	tagcagagcg	aggatgttag	gcggtgctac	2340
agagttcttg	aagtgggtggc	ctaactacgg	ctacactaga	aggacagtat	ttggtatctg	2400
cgctctgctg	aagccagtta	ccttcggaaa	aagagttggt	agctcttgat	ccggcaaaca	2460
aaccaccgct	ggtagcggtg	gtttttttgt	ttgcaagcag	cagattacgc	gcagaaaaaa	2520
aggatctcaa	gaagatcctt	tgatcttttc	tacggggtct	gacgctcagt	ggaacgaaaa	2580
ctcacgttaa	gggatttttg	tcattgagatt	atcaaaaagg	atcttcacct	agatcctttt	2640
aaattaaaaa	tgaagtttta	aatcaatcta	aagtatatat	gagtaaaact	ggtctgacag	2700
ttaccaatgc	ttaatcagtg	aggcacctat	ctcagcgatc	tgtctatttc	gttcatccat	2760
agttgcctga	ctccccgtcg	tgtagataac	tacgatacgg	gagggcttac	catctggccc	2820
cagtgtgcga	atgataccgc	gagacccacg	ctcaccggct	ccagatttat	cagcaataaa	2880
ccagccagcc	ggaagggccg	agcgagaaag	tggtcctgca	actttatccg	cctccatcca	2940
gtctattaat	tggtgccggg	aagctagagt	aagtagttcg	ccagttaata	gtttgcgcaa	3000
cgttggtggc	attgctacag	gcacgtggtg	gtcacgctcg	tcgtttggta	tggettcat	3060
cagctccggt	tccaacgat	caaggcgagt	tacatgatcc	cccatgttgt	gcaaaaaagc	3120
ggttagctcc	ttcggtcctc	cgatcgttgt	cagaagtaag	ttggccgcag	tggtatcact	3180
catgggttat	gcagcactgc	ataattctct	tactgtcatg	ccatccgtaa	gatgcttttc	3240
tgtgactggt	gagtactcaa	ccaagtcatt	ctgagaatac	cgcgcccggc	gaccgagttg	3300
ctcttgcccg	gcgtcaatac	gggataatag	tgtatgacat	agcagaactt	taaaagtgt	3360

```

catcattgga aaacgttctt cggggcgaaa actctcaagg atcttaccgc tgttgagatc 3420
cagttcgatg taaccactc gtgcacccaa ctgatcttca gcatctttta ctttcaccag 3480
cgtttctggg tgagcaaaaa caggaaggca aaatgccgca aaaaaggga taagggcgac 3540
acggaaatgt tgaatactca tactcttctt ttttcaatat tattgaagca tttatcaggg 3600
ttattgtctc atgagcggat acatatttga atgtatttag aaaaataaac aaataggggt 3660
tccgcgcaca tttccccgaa aagtgccacc tgacgtctaa gaaaccatta ttatcatgac 3720
attaacctat aaaaataggc gtatcacgag gccctttcgt ctgcgcggtt tcggtgatga 3780
cggtgaaaac ctctgacaca tgcagctccc ggagacggtc acagcttgtc tgtaagcggg 3840
tgccgggagc agacaagccc gtcagggcgc gtcagcgggt gttggcgggt gtcggggctg 3900
gcttaactat gcggcatcag agcagattgt actgagagtg caccatatgc ggtgtgaaat 3960
accgcacaga tgcgtaagga gaaaataccg catcaggcga aattgtaaac gttaatattt 4020
tgttaaaatt cgcgttaaat atttgttaaa tcagctcatt ttttaacca taggccgaaa 4080
tcggcaaaat cccttataaa tcaaaagaat agaccgagat aggggtgagt gttgttccag 4140
tttgaacaa gagtccacta ttaaagaacg tggactcaa cgtcaaagg cgaaaaaccg 4200
tctatcaggg cgatggccca ctacgtgaac catcacccaa atcaagtttt ttgcggtcga 4260
ggtgccgtaa agctctaaat cggaacccta aaggagagccc ccgatttaga gcttgacggg 4320
gaaagccggc gaacgtggcg agaaaggaag ggaagaaagc gaaaggagcg ggcgctaggg 4380
cgctggcaag tgtagcggtc acgctgcgcg taaccaccac acccgccgcg cttaatgcgc 4440
cgctacaggg cgcgtccatt cgccattcag gctgcgcaac tgttggaag ggcgatcgg 4500
gcgggcctct tcgctattac gccagctggc gaaaggggga tgtgctgcaa ggcgattaag 4560
ttgggtaacg ccagggtttt ccagtcacg acgttgtaaa acgacggcca gtgaattgta 4620
atacgactca ctatagggcg aattcaaaaa accctcaag acccgtttag agggcccaag 4680
gggttatgct agtgaattct gcagggtacc cggggatcct ctagagatcc ctcgacctcg 4740
agatccattg tgctgg 4756

```

<210> 4

<211> 4643

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX52

<400> 4

```

gagtgcacca tatgcgggtg gaaataccgc acagatgcgt aaggagaaaa taccgcatca 60
ggcgaaattg taaacgttaa tattttgtta aaattcgctg taaatatttg ttaaatcagc 120
tcatttttta accaataggc cgaaatcggc aaaatccctt ataaatcaaa agaatagacc 180
gagatagggt tgagtgttgt tccagtttgg aacaagagtc cactattaaa gaacgtggac 240
tccaacgtca aagggcgaaa aaccgtctat cagggcgatg gccactacg tgaacctca 300
cccaaataca gttttttgcg gtcgaggtgc cgtaaagctc taaatcggaa ccctaaagg 360
agcccccgat ttagagcttg acgggggaaag ccggcgaaag tggcgagaaa ggaagggaag 420
aaagcgaaag gagcggggcg tagggcgctg gcaagtgtag cggtcacgct gcgcgtaacc 480
accacaccgg ccgcgcttaa tgcgccgcta cagggcgctg ccattcgcca ttcaggctgc 540
gcaactgttg ggaaggcgga tcggtgcggg cctcttcgct attacgccag ctggcgaaag 600
gggatgtgc tgcaaggcga ttaagttggg taacgccagg gttttccag tcacgacgtt 660
gtaaaacgac ggccagtga ttgtaatacg actcactata gggcgaaatt aaaaaacccc 720
tcaagaccgg tttagaggcc ccaaggggtt atgctagtga attctgcagg gtaccgggg 780
atcctctaga gatccctcga cctcgagatc cattgtgctg gcagccgatc tccgtcttgt 840
gaagatctac tccaccacca tccgtatcga tcagtccatc ctcaccggag aatctgtgtc 900
tgttatcaag cacaccgact ctgtgccaga tccacgcgct gtttaaccag acaagaagaa 960
ttgtctgttc tcgggaacca atgtcgcac tggaaaggct cgtggaatcg tcttcggaac 1020
cggattgacc actgaaatcg gaaagatccg taccgaaatg gctgagaccg agaatgagaa 1080
gacaccactt caacagaagt tggacgaatt cggagagcaa ctttccaagg ttatctctgt 1140
tatttgcgtt gctgtttggg ctatcaacat tggacatttc aacgatccag ctcacgggtg 1200
atcatgggtt aaggagcaa tctactactt caaaatcgcc gttgctcttg ccgtcgctgc 1260
tattccagaa ggacttccag ctgtcatcac cacgtgcctt gccctcgga ctcgcgctat 1320
ggccaagaag aacgctattg taagatccct tccatccgtc gaaactcttg gatgcacatc 1380
tgttatctgc tctgacaaga ctggaactct caccaccaac cagatgtctg tgtcaaagat 1440

```

gttcacgcgt	ggacaagctt	ctggagacaa	catcaacttc	accgagttcg	ccatctccgg	1500
atccacctac	gagccagtcg	gaaaggtttc	caccaatgga	cgtgaaatca	acccagctgc	1560
tggagaatte	gaatcactca	ccgagttggc	catgatctgc	gctatgtgca	atgattcatc	1620
tgttgattac	aatgagacca	agaagatcta	cgagaaagtc	ggagaagcca	ctgaaactgc	1680
tcttatcggt	cttgctgaga	agatgaatgt	tttcggaacc	tcgaaagccg	gactttcacc	1740
aaaggagctc	ggaggagttt	gcaaccgtgt	catccaacaa	aatggaaga	aggagttcac	1800
actcgagttc	tcccgtgatc	gtaaatccat	gtccgcctac	tgcttcccag	cttccggagg	1860
atctggagcc	aagatgttcg	tgaaggagc	cccagaagga	gttctcggaa	gatgcaccca	1920
cgtcagagtt	aacggacaaa	aggttccact	cacctctgcc	atgactcaga	agattgttga	1980
ccaatgctgt	caatacggaa	ccggaagaga	tacccttcgt	tgtcttgccc	tcggccagca	2040
caatggatct	cgagggatct	tccataccta	ccagttctgc	gcctgcaggt	cgcgcccgcg	2100
actctctaga	cgcgtaagct	tactagcata	accccttggg	gcctctaaac	gggtcttgag	2160
gggttttttg	agcttctcgc	cctatagtga	gtcgtattac	agcttgagta	ttctatagt	2220
tcacctaaat	agcttgcgct	aatcatggtc	atagctgttt	cctgtgtgaa	attgttatcc	2280
gctcacaatt	ccacacaaca	tacgagccgg	aagcataaag	tgtaaagcct	ggggtgccta	2340
atgagtgage	taactcacat	taattgcgtt	gcgctcactg	cccgttttcc	agtcgggaaa	2400
cctgtcgtgc	cagctgcatt	aatgaatcgg	ccaacgcgcg	gggagaggcg	gtttgcgtat	2460
tgggcgctct	tccgcttcc	cgctcactga	ctcgtcgcgc	tcggtcgttc	ggctgcggcg	2520
agcggtatca	gctcactcaa	aggcggtaat	acggttatcc	acagaatcag	gggataacgc	2580
aggaaagaac	atgtgagcaa	aaggccagca	aaaggccagg	aaccgtaaaa	aggccgcgtt	2640
gctggcggtt	ttcgataggc	tccgcccccc	tgacgagcat	cacaaaaatc	gacgctcaag	2700
tcagagggtg	cgaaacccga	caggactata	aagataccag	gcgtttcccc	ctggaagctc	2760
cctcgtgcgc	tctcctgttc	cgaccctgcc	gcttaccgga	tacctgtccg	cctttctccc	2820
ttcgggaagc	gtggcgcttt	ctcatagctc	acgctgtagg	tatctcagtt	cggtgtaggt	2880
cgttcgctcc	aagctgggct	gtgtgcacga	acccccggtt	cagcccgacc	gctgcgcctt	2940
atccggtaac	tatcgtcttg	agtccaaccc	ggtaagacac	gacttatcgc	cactggcagc	3000
agccactggt	aacaggatta	gcagagcgag	gtatgtaggc	ggtgctacag	agttcttgaa	3060
gtggtggcct	aactacggct	acactagaag	gacagtattt	ggtatctgcg	ctctgctgaa	3120
gccagttacc	ttcggaaaaa	gagttggtag	ctcttgatcc	ggcaaacaaa	ccaccgctgg	3180
tagcggtggt	ttttttgttt	gcaagcagca	gattacgcgc	agaaaaaaag	gatctcaaga	3240
agatcctttg	atcttttcta	cggggtctga	cgctcagtgg	aacgaaaact	cacgttaagg	3300
gatttttggtc	atgagattat	caaaaaggat	cttcacctag	atccttttaa	attaaaaatg	3360
aagtttttaa	tcaatctaaa	gtatatatga	gtaaacttgg	tctgacagtt	accaatgctt	3420
aatcagtgag	gcacctatct	cagcgatctg	tctatttcgt	tcatccatag	ttgcttgact	3480
cccgcgtcgt	tagataacta	cgatacggga	gggcttacca	tctggcccca	gtgctgcaat	3540
gataccgcga	gacccacgct	caccggctcc	agatttatca	gcaataaacc	agccagccgg	3600
aagggccgag	cgcagaagtg	gtcctgcaac	tttatccgcc	tccatccagt	ctattaattg	3660
ttgcccggaa	gctagagtaa	gtagtccgcc	agttaatagt	ttgcgcaacg	ttgttgcat	3720
tgctacaggc	atcgtggtgt	cacgctcgtc	gtttggtatg	gcttcattca	gctccggttc	3780
ccaacgatca	aggcgagtta	catgatcccc	catgttgtgc	aaaaaagcgg	ttagctcctt	3840
cggctcctcc	atcgttgtca	gaagtaagtt	ggccgcagtg	ttatcactca	tggttatggc	3900
agcactgcat	aattctctta	ctgtcatgcc	atccgtaaga	tgcttttctg	tgactggtga	3960
gtactcaacc	aagtcattct	gagaataccg	cgcccggcga	ccgagttgct	cttgcccggc	4020
gtcaatacgg	gataatagt	tatgacatag	cagaacttta	aaagtgtca	tcattggaaa	4080
acgttcttcg	gggcgaaaac	tctcaaggat	cttaccgctg	ttgagatcca	gttcgatgta	4140
acccactcgt	gcacccaact	gatcttcagc	atcttttact	ttcaccagcg	tttctgggtg	4200
agcaaaaaca	ggaaggcaaa	atgccgcaaa	aaagggaata	agggcgacac	ggaaatgttg	4260
aatactcata	ctcttccttt	ttcaatatta	ttgaagcatt	tatcagggtt	attgtctcat	4320
gagcggatac	atatttgaat	gtatttagaa	aaataaacia	ataggggttc	cgcgcacatt	4380
tccccgaaaa	gtgccacctg	acgtctaaga	aaccattatt	atcatgacat	taacctataa	4440
aaataggcgt	atcacgaggc	cctttcgtct	cgcgcgtttc	ggtgatgacg	gtgaaaacct	4500
ctgacacatg	cagctcccgg	agacggtcac	agcttgtctg	taagcggatg	ccgggagcag	4560
acaagcccgt	cagggcgcggt	cagcgggtgt	tggcgggtgt	cggggcgtggc	ttaactatgc	4620
ggcatcagag	cagattgtac	tga				4643

<210> 5
 <211> 4454
 <212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGX104

<400> 5

gagtgcacca	tatgcggtgt	gaaataccgc	acagatgcgt	aaggagaaaa	taccgcatca	60
ggcgaaattg	taaacgttaa	tattttgtta	aaattcgcgt	taaatatttg	ttaaatacgc	120
tcatttttta	accaataggc	cgaaatcggc	aaaatccctt	ataaatcaaa	agaatagacc	180
gagataggg	tgagtgttgt	tccagtttgg	aacaagagtc	cactattaaa	gaacgtggac	240
tccaacgtca	aagggcgaaa	aaccgtctat	cagggcgatg	gcccactacg	tgaaccatca	300
cccaaataca	gttttttgcg	gtcagaggtg	cgtaaagctc	taaatcggaa	ccctaaagg	360
agccccgat	ttagagcttg	acggggaaa	ccggcgaaac	tggcgagaaa	ggaagggaag	420
aaagcgaaa	gagcggggcg	tagggcgctg	gcaagtgtag	cggtcacgct	gcgcgtaacc	480
accacacccg	ccgcgcttaa	tgcgccgcta	cagggcgcg	ccattcgcca	ttcaggctgc	540
gcaactgttg	ggaaggcgga	tccgtgcggg	cctcttcgct	attacgccag	ctggcgaaa	600
ggggatgtgc	tgcaaggcga	ttaagttggg	taacgccagg	gttttcccag	tcacgacgtt	660
gtaaaacgac	ggccagtga	ttgtaatacg	actcactata	gggcgaattc	aaaaaacccc	720
tcaagacccg	tttagaggcc	ccaaggggtt	atgctagtga	attctgcagg	gtacccgggg	780
atcctctaga	gatccctcga	cctcgagatc	cattgtgctg	gaccgtggta	ctcttatgga	840
gctcggaatc	tcgccaatcg	tcacttctgg	acttatcatg	caacttctcg	ccggagccaa	900
gatcatcgaa	gtcggagaca	caccaaagga	ccgtgctctt	ttcaacggag	cccagaaatg	960
taagccgaaa	agtgtgtgtt	ttcaatctct	aatttttgaa	cttttcagtg	ttcggtatgg	1020
tcactactgt	tggaacagct	attgtctacg	tcattgtcgg	actctacgga	gagccatcgg	1080
aaatcggagc	tggaatctgt	ctccttatcg	tcgtccaact	cggtattgcc	ggtctcatcg	1140
tcctccttct	cgacgagctt	ctccaaaagg	gatatgggtc	cggatccgga	atttctctct	1200
tcattgccac	caacatctgt	gaaaccattg	tctggaaggc	attctccccg	gcaacaatga	1260
acaccggacg	tggaaccgag	ttcgaaggag	ccgtcattgc	tcttttccat	cttcttgcca	1320
cccgtccga	caaggtccgt	gcccttcgtg	aggctttcta	ccgtcaaaac	cttccaaact	1380
tgatgaactt	gatggctact	ttcctcgttt	ttgcgggtgt	tatctacttc	caaggattcc	1440
gtgtcgacct	cccaatcaag	tctgcccgtc	accgtggaca	atacagcagc	taccaaatca	1500
agctcttcta	cacctccaac	attccaatca	tccttcaatc	tgctctcgtc	tccaacctct	1560
acgttatctc	tcaggtttgt	tgcattctcag	tagtaccgtt	agatgtttat	ctttctctag	1620
aggggtcaagt	tggecgagaa	attttttgag	ttcattctca	agtctgatgg	aaaatgttta	1680
tttttcagat	gtcgcgcgga	aagttcggag	gaaacttctt	catcaacctt	ctcggtacct	1740
ggtecgataa	caccggatac	agaagctacc	caactggagg	actctgctac	tatctttcac	1800
caccagagtc	tcttggaac	atcttcgaag	acccaatcca	ctgcaccagc	acaatggatc	1860
tcgagggatc	ttccatacct	accagttctg	cgcttcgagg	tcgcggccgc	gactctctag	1920
acgcgtaagc	ttactagcat	aaccccttgg	ggcctctaaa	cgggtcttga	gggggttttt	1980
gagcttctcg	ccctatagtg	agtcgtatta	cagcttgagt	attctatagt	gtcacctaaa	2040
tagcttggcg	taatcatggt	catagctgtt	tcctgtgtga	aattgttatc	cgctcacaat	2100
tccacacaac	atacagaccg	gaagcataaa	gtgtaaagcc	tggggtgcct	aatgagttag	2160
ctaactcaca	ttaattgcgt	tgcgctcact	gcccgctttc	cagtcgggaa	acctgtcgtg	2220
ccagctgcat	taatgaatcg	gccaacgcgc	ggggagaggc	ggtttgcgta	ttgggcgctc	2280
ttccgcttcc	tcgctcactg	actcgctgcg	ctcggtcggt	cggctgcggc	gagcgggtatc	2340
agctcactca	aaggcggtaa	tacggttatc	cacagaatca	ggggataacg	caggaaagaa	2400
catgtgagca	aaaggccagc	aaaaggccag	gaaccgtaaa	aaggccgcgt	tgctggcggt	2460
tttcgatagg	ctccgcccc	ctgacgagca	tcacaaaaat	cgacgctcaa	gtcagagggtg	2520
gcgaaacccg	acaggactat	aaagatacca	ggcgtttccc	cctggaagct	ccctcgtgcg	2580
ctctcctggt	ccgacctgc	cgcttaccgg	atacctgtcc	gcctttctcc	cttcgggaag	2640
cgtggcgctt	tctcatagct	cacgctgtag	gtatctcagt	tcgggtgtagg	tcgttcgctc	2700
caagctgggc	tgtgtgcacg	aacccccgt	tcagcccagc	cgctgcgcct	tatccggtaa	2760
ctatcgtctt	gagtccaacc	cggtaaagca	cgacttatcg	ccactggcag	cagccactgg	2820
taacaggatt	agcagagcga	ggtatgtagg	cggtgctaca	gagttcttga	agtgggtggc	2880
taactacggc	tacactagaa	ggacagtatt	tggtatctgc	gctctgctga	agccagttac	2940
cttcggaaaa	agagttggta	gctcttgatc	cggcaaaaaa	accaccgctg	gtagcgggtg	3000
tttttttgtt	tgcaagcagc	agattacgcg	cagaaaaaaa	ggatctcaag	aagatccttt	3060
gatcttttct	acggggtctg	acgctcagtg	gaacgaaaaa	tcacgttaag	ggatttttgt	3120


```
catgagatta tcaaaaagga tcttcaccta gatcctttta aattaaat gaagtttta 3180
atcaatctaa agtatatatg agtaaacttg gtctgacagt taccaatgct taatcagtga 3240
ggcacctatc tcagcgatct gtctatttctg ttcacccata gttgcctgac tccccgtcgt 3300
gtagataact acgatacggg agggcttacc atctggcccc agtgctgcaa tgataccgcg 3360
agaccacgc tcaccggctc cagatttatc agcaataaac cagccagccg gaagggccga 3420
gcgcagaagt ggtcctgcaa ctttatccgc ctccatccag tctattaatt gttgccggga 3480
agctagagta agtagttcgc cagttaatag tttgcgcaac gttgttgga ttgctacagg 3540
catcgtggtg tcacgctcgt cgtttggtat ggcttcattc agctccggtt cccaacgatc 3600
aaggcgagtt acatgatccc ccatgttggtg caaaaagcg gttagctcct tcggtcctcc 3660
gatcgttgtc agaagtaagt tggccgcagt gttatcactc atggttatgg cagcactgca 3720
taattctctt actgtcatgc catccgtaag atgcttttct gtgactggtg agtactcaac 3780
caagtcattc tgagaatacc gcgcccggcg accgagttgc tcttgcccg cgtcaatacg 3840
ggataatagt gtatgacata gcagaacttt aaaagtgtc atcattggaa aacgttcttc 3900
ggggcgaaaa ctctcaagga tcttaccgct gttgagatcc agttcgatgt aaccactcg 3960
tgcacccaac tgatcttcag catcttttac tttcaccagc gtttctgggt gagcaaaaac 4020
aggaaggcaa aatgccgcaa aaaagggaat aaggcgaca cggaaatgtt gaatactcat 4080
actcttcctt tttcaatatt attgaagcat ttatcagggt tattgtctca tgagcggata 4140
catatttgaa tgtatttaga aaaataaaca aataggggtt ccgcgcacat tccccgaaa 4200
agtgccacct gacgtctaag aaaccattat tatcatgaca ttaacctata aaaataggcg 4260
tatcacgagg ccctttcgtc tcgcgcgttt cggtgatgac ggtgaaaacc tctgacacat 4320
gcagctccog gagacggtca cagcttgtct gtaagcggat gccgggagca gacaagccc 4380
tcagggcgcg tcagcgggtg ttggcgggtg tcggggctgg cttaactatg cggcatcaga 4440
gcagattgta ctga 4454
```

<210> 6

<211> 4701

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence: Plasmid pGZ18

<400> 6

```
accagcttt cttgtacaaa gtggtgatct ttccagcaca atggatctcg agggatcttc 60
catacctacc agttctgctc ctgcaggtcg cggccgcgac tctctagacg cgtaagctta 120
ctagcataac cccttggggc ctctaaacgg gtcttgaggg gttttttgag cttctcgccc 180
tatagttagt cgtattacag cttgagtatt ctatagtgtc acctaaatag cttggcgtaa 240
tcatggtcat agctgtttcc tgtgtgaaat tgttatccgc tcacaattcc acacaacata 300
cgagccggaa gcataaagtg taaagcctgg ggtgcctaata gagtgagcta actcacatta 360
attgcgttgc gctcactgcc cgctttccag tcgggaaacc tgtcgtgcca gctgcattaa 420
tgaatcggcc aacgcgcggg gagaggcggg ttgctgattg ggcgctcttc cgcttcctcg 480
ctcactgact cgctgcgctc ggtcgttcgg ctgcggcgag cggtatcagc tcaactcaaag 540
gcggtaatac ggttatccac agaatcaggg gataacgcag gaaagaacat gtgagcaaaa 600
ggccagcaaa aggccaggaa ccgtaaaaag gccgcgttgc tggcgttttt cgataggctc 660
cgccccctg acgagcatca caaaaatcga cgctcaagtc agaggtggcg aaaccogaca 720
ggactataaa gataccaggc gtttccccct ggaagctccc tcgtgcgctc tcctgttccg 780
accctgccgc ttaccggata cctgtccgcc tttctccctt cgggaagcgt ggcgctttct 840
catagctcac gctgtaggta tctcagttcg gtgtaggctg ttcgctccaa gctgggctgt 900
gtgcacgaac ccccggttca gcccgaccgc tgcgccttat ccggttaacta tcgtcttgag 960
tccaaccggg taagacacga cttatcgcca ctggcagcag ccactggtaa caggattagc 1020
agagcgaggt atgtaggcgg tgctacagag ttcttgaagt ggtggcctaa ctacggctac 1080
actagaagga cagtatttgg tatctgcgct ctgctgaagc cagttacctt cggaaaaaga 1140
gttggttagct cttgatccgg caaacaacc accgctggtg gcggtggttt ttttgtttgc 1200
aagcagcaga ttacgcgcag aaaaaagga tctcaagaag atcctttgat cttttctacg 1260
gggtctgacg ctacgtggaa cgaaaactca cgttaaggga ttttgggtcat gagattatca 1320
aaaaggatct tcacctagat ctttttaaat taaaaatgaa gtttttaaat aatctaaagt 1380
atatatgagt aaacttggtc tgacagttac caatgcttaa tcagtaggc acctatctca 1440
gcgatctgtc tatttcgttc atccatagtt gcctgactcc ccgtcgtgta gataactacg 1500
```

atacgggagg gcttaccatc tggccccagt gctgcaatga taccgcgaga cccacgctca 1560
ccggctccag atttatcagc aataaaccag ccagccggaa gggccgagcg cagaagtggg 1620
cctgcaactt tatccgcctc catccagtct attaattgtt gccgggaagc tagagtaagt 1680
agttcgccag ttaatagttt gcgcaacggt gttggcattg ctacaggcat cgtggtgtca 1740
cgctcgtcgt ttggtatggc ttcattcagc tccggttccc aacgatcaag gcgagttaca 1800
tgatccccc tgttgtgcaa aaaagcgggt agctccttcg gtcctccgat cgttgtcaga 1860
agtaagtgtg ccgcagtgtt atcactcatg gttatggcag cactgcataa ttctcttact 1920
gtcatgccat ccgtaagatg cttttctgtg actggtgagt actcaaccaa gtcattctga 1980
gaataccgcg cccggcgacc gagttgctct tgcccggcgt caatacggga taatagtgt 2040
tgacatagca gaactttaaa agtgctcatc attgaaaaac gttcttcggg gcgaaaactc 2100
tcaaggatct taccgctgtt gagatccagt tcgatgtaac ccactcgtgc acccaactga 2160
tcttcagcat cttttacttt caccagcgtt tctgggtgag caaaaacagg aaggcaaaat 2220
gccgcaaaaa agggaataag ggcgacacgg aaatgttgaa tactcatact cttccttttt 2280
caatattatt gaagcattta tcagggttat tgtctcatga gcggatacat atttgaatgt 2340
atttagaaaa ataaacaaat aggggttccg cgcacatttc cccgaaaagt gccacctgac 2400
gtctaagaaa ccattattat catgacatta acctataaaa ataggcgtat cagcagggccc 2460
tttcgtctcg cgcgtttcgg tgatgacggt agcggatgcc gggagcagac aagcccgtca gggcgcgtca 2520
acggtcacag cttgtctgta agcggatgcc gggagcagac aactatgcgg catcagagca gattgtactg 2580
gcgggtgttg gcgggtgtcg gggctggctt aactatgcgg catcagagca gattgtactg 2640
agagtgcacc atatgcggtg tgaaataccg cacagatgcg taaggagaaa ataccgcatc 2700
aggcgaaatt gtaaacgtta atattttgtt aaaattcgcg ttaaataatt gttaaatacag 2760
ctcatttttt aaccaatagg ccgaaatcgg caaaatccct tataaatcaa agaataagac 2820
cgagataggg ttgagtgttg ttccagtttg gaacaagagt ccactattaa agaacgtgga 2880
ctccaacgtc aaaggcgcaa aaaccgtcta tcagggcgat ggcccactac gtgaaccatc 2940
acccaaatca agttttttgc ggtcgaggtg ccgtaaagct ctaaatacga accctaaagg 3000
gagccccga ttttagagctt gacggggaaa gccggcgaaac gtggcgagaa aggaaggga 3060
gaaagcgaaa ggagcggcg ctaggcgct ggcaagtgt gcggtcacgc tgcgcgtaac 3120
caccacaccc gccgcgctta atgcgcgct acaggcgcg gctcttcgc tattacgcca gctggcgaaa 3180
cgcaactgtt gggaaggcg atcgggtgcg gctcttcgc tattacgcca gctggcgaaa 3240
gggggatgt ctgcaaggcg attaagtttg gtaacgccag ggttttccca gtcacgacgt 3300
tgtaaaacga cggccagtga attgtaatac gactcactat agggcgaaat caaaaaaccc 3360
ctcaagaccc gtttagaggg cccaaggggt tatgctagt ccattgtgct ggaaagcctt tgcagggtg 3420
gatectctag agatccctcg acctcgagat ccattgtgct ggaaagcctt tgcagggtg 3480
gcaagccacg tttggtggtg gcgaccatcc tccaaaatca acaagtttgt acaaaaaagc 3540
aggctatgcc aagtacatgt cgattgcgta cgcgttcgta atgttggctg tgttagtcgc 3600
taccagcagt caaattgttc tcgagagtgc gtttttacat tatcccttca tcttgattac 3660
gacaattttc agctgttctc gtcctacat ctctcttcat tgtcacaatg gtcggaatct 3720
tcttctttgc tgcattgtct catccaaaag aattcacgaa tattatccat ggtgtcgtat 3780
tcttctcat gattccatct acatatgtgt tctcacttt atattcgtc atcaatctca 3840
acgttatcac gtggggaact cgtgaagctg tcgctaaggc aacgggacaa aagacgaaaa 3900
aagcgcctat ggaacaattt atagacagag tgattgatat tgtgaaaaag ggattcagat 3960
taatcagttg tcgggagaag aaggaacatg aagagagac agagaaaatg gaaaagaaaa 4020
tgacagagaat ggagctagcc ttgagaagta ttgaggttat ctttaacttt agaaatgtga 4080
aattaataat ttattttcag agtgggtgcc acgtgaagaa aattctcgat gcaacagagg 4140
agaaggagaa acgtgaagaa gaaactcaaa ctgcagattt tccgattgaa gagaacgtag 4200
agaagactca aaaagagatt cagaaggcaa accgttatgt gtggatgaca agtcatagct 4260
tgaaagtgtg tgaacgagga aaactgaaaa gtgcggaaaa ggttttctgg aacgagctca 4320
tcaatgcata tctgaaaccg atcaagacga cgccagctga aatgaaagcc gtcgccgaag 4380
gattggcttc tctacgaaat cagattgctt tcactattct tctcgttaat tctcttcttg 4440
ctcttgccat ctttttgatt cagaaacaca aaaatgtgct cagcatcaag ttctcgccaa 4500
tcagtaagca atattacctt tatggtcaat tcaaaaaatt tgtttttttt ttctagaaaa 4560
cttccgatgg acgaaaatga atgagatgac tggacaatac gaggaaccg atgaaccatt 4620
aaaaatagat ccacttgga tgggaattgt tgttttcctt ctaattattc tttttgttca 4680
aactctcgga atgcttctcc a 4701

<210> 7

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:
Oligonucleotide primer C04H5.6F

<400> 7

tgctcagaga gtttctcaac gaacc

25

<210> 8

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer C04H5.6R

<400> 8

caatgtagt tgctaggacc acctg

25

<210> 9

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bF

<400> 9

cagccgatct ccgtcttgtg

20

<210> 10

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer K11D9.2bR

<400> 10

ccgagggcaa gacaacgaag

20

<210> 11

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15F

<400> 11

accgtggtac tcttatggag ctcg

24

<210> 12

<211> 20
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer Y57G11C.15R

<400> 12
tgcagtggat tgggtcttcg 20

<210> 13
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2F

<400> 13
ggggacaagt ttgtacaaaa aagcaggcta tgccaagtac atgtcgattg cg 52

<210> 14
<211> 52
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:Oligonucleotide
primer T25G3.2R

<400> 14
ggggaccact ttgtacaaga aagctggggtt ggagaagcat tccgagagtt tg 52